

## 53 BROADLEAVES. 11 GRASSES. ONE JUG.


Growers now have a better and more convenient way to control weeds in spring, durum and winter wheat. New BATALIUM™ Herbicide is driven by a unique, high-performance formulation with three modes of action that provide all-in-one weed control in a single application. With BATALIUM, you'll get long-lasting control of your toughest grass weeds like wild oats and green foxtail, in addition to dependable, wide-spectrum control of broadleaf weeds like common lambsquarters, wild buckwheat and more.

### FEATURES & BENEFITS

- » Advanced formulation of flucarbazone-sodium, a highly effective Group 2 herbicide, with fluroxypyr (Group 4) and bromoxynil (Group 6)
- » Broad-spectrum broadleaf weed control
- » Long-lasting control of hard-to-kill grassy weeds
- » Best-in-class crop safety
- » Compatible with multiple tank-mix partners


## UNMATCHED WEED CONTROL

AVERAGE WEED CONTROL 5 WEEKS AFTER APPLICATION



## BEST-IN-CLASS CROP SAFETY

AVERAGE CROP INJURY TO SPRING WHEAT 2 WEEKS AFTER APPLICATION



## TIMING AND USAGE RATES

### CROP TIMING

Apply from 2-leaf stage up to 60 days prior to harvest. Winter wheat applications can be made in the fall or spring.

### WEED TIMING

Make application to control grass weeds 1-4 leaf stage. Make application to control broadleaf weeds up to 4 inches tall.

### RATE

BATALIUM has a convenient single use rate of 13.7 fl oz/A.

### TANK-MIX FLEXIBILITY

Compatible with over 23 registered tank-mix partners, BATALIUM lets you customize your weed management program based on weed pressure and application conditions.

## WEEDS CONTROLLED

GRASSES	BROADLEAVES <sup>1</sup>		
Green foxtail	Annual sowthistle	Kochia <sup>2</sup>	Ragweed (common, giant)
Volunteer oat	Bristly starbur	Lambsquarters, common	Russian thistle
Wild oat	Buffalobur	Lanceleaf sage	Shepherd's-purse
Yellow foxtail <sup>4</sup>	Burcucumber	London rocket	Smartweed (green, ladythumb, Pennsylvania)
Japanese brome <sup>P</sup>	Catchweed bedstraw (cleavers, 1-4 whorls)	Mallow (common, Venice)	Sunflower <sup>3</sup>
True cheat <sup>P</sup>	Coast fiddleneck	Morning glory (ivyleaf, pitted, tall)	Tartary buckwheat
Barnyard grass <sup>P4</sup>	Cocklebur, common	Mustard (blue, wild)	Tarweed, common
Downy brome <sup>P4</sup>	Corn gromwell	Nightshade (black, cutleaf, Eastern black, hairy, silverleaf)	Velvetleaf
Foxtail barley <sup>P4</sup>	Cow cockle	Pepperweed	Volunteer canola
Persian darnel <sup>P4</sup>	Field pennycress, field	Pigweed (redroot, spiny)	Volunteer flax
Rattail fescue <sup>P4</sup>	Grounsel, common	Prickly lettuce	Wild buckwheat
	Hemp sesbania	Puncturevine	Wild radish
	Jimsonweed	Purslane, common	Yellow starthistle
	Knawel		
	Knotweed		

<sup>P</sup> Partially controlled for grasses.

<sup>1</sup> Up to 4 inches in height; 2-inch rosette. For the states of Idaho, Oregon, Washington, Colorado, Wyoming, and Montana, add bromoxynil to provide 0.125 to 0.25 lb ai/A. Target weed size up to 2 inches in height; 1 inch rosette.

<sup>2</sup> Includes herbicide tolerant or resistant. Best control is achieved when weeds are at least 1 inch tall.

<sup>3</sup> For best control of sunflower, delay application until emerging seedlings are 4 inches in height.

<sup>4</sup> A tank-mix with AUDIT® brand herbicides or other herbicides containing tribenuron is required to achieve the level of control listed above.



**Always read and follow label directions.** AUDIT is a registered trademark and BATALIUM is a trademark of an Arysta LifeScience Group Company. Huskie is a registered trademark of Bayer CropScience. PerfectMatch is a registered trademark of Dow AgroSciences LLC. Axial and Talinor are registered trademarks of a Syngenta Group Company. Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation. ©2018 Arysta LifeScience Group Company. BATU-006